



Technical Information Sheet
Article No. 9067

H-F Antidote Gel

1 component gel for the treatment of hydrofluoric acid burns

Range of use

For the first aid treatment of burns to the skin, caused by Hydrofluoric Acid.

Application

Initially wash the affected area with copious amounts of water for at least 1 minute. Carefully remove contaminated clothing, taking care not to contaminate further areas of the skin. Apply H-F Antidote Gel on and around the affected area with gentle massage. Seek medical attention. Continue to massage the gel into the affected area, using repeated applications until 15 minutes after the pain has subsided, or until medical treatment is

Characteristic data of the product

Active ingredient	Calcium Gluconate 2.5% w/w
-------------------	----------------------------

available. If after medical treatment, pain should reoccur at the burn site re-apply the H-F Antidote Gel and seek further medical attention immediately.

Discard all opened tubes of H-F Antidote Gel after treatment.

Packaging

25g aluminium tube

Shelf life

30 months in unopened packaging in a cool and dry storage place at temperatures between +8°C and +25°C.

The statements above are compiled from our field of production and according to the latest technological developments and application techniques.

Since application and working are beyond our control, no liability of the producer can be derived from the contents of this information sheet. Any statements made beyond the contents of this information must be confirmed in writing by the producer.

In all cases, our general conditions of sale are valid. With the publication of this Technical Information Sheet all previous editions are no longer valid.



Remmers (UK) Limited Crawley
United Kingdom
Tel: +44(0) 845 373 0103
Fax: +44(0)845 373 0104
www.remmers.co.uk

Remmers (Far East) Pte. Ltd.
Singapore
Tel: +65 6 7410277
Fax: +65 6 7417158